

Invited: QMBS-I-06

Screened exchange dynamical mean field theory

Silke Biermann

Centre de Physique Theorique, Ecole Polytechnique, France

Dynamical mean field theory (DMFT), in conjunction with electronic structure techniques such as density functional theory (DFT), has led to tremendous progress in the description of excited state properties of materials with strong electronic Coulomb correlations. The challenge nowadays consists in refining the interface of electronic structure and many-body theory in order to develop quantitatively accurate predictive schemes.

We review recent efforts of incorporating dynamical screening effects into a DMFT-based description of correlated materials [1]. Such effects can stem either from higher energy degrees of freedom that have been integrated out [2] or from nonlocal processes that are effectively backfolded into a local description. This can be conveniently done by combined many-body perturbation theory and dynamical mean field theory ("GW+DMFT") techniques [3]. These insights lead to a simple but efficient scheme, dubbed "Screened exchange dynamical mean field theory" [4], which can be understood as an approximation to GW+DMFT, or as a non-perturbative dynamical generalization of Hedin's Coulomb-hole-screened-exchange approximation. In particular, it includes non-local exchange beyond the local density approximation and dynamical screening beyond standard DMFT techniques.

References:

10. S. Biermann, *J. Phys.: Condens. Matter*, **26**, 173202 (2014).
2. J.M. Tomczak et al., *Phys. Rev. B*, **90**, 165138 (2014).
3. T. Ayral et al., *Phys. Rev. B*, **87**, 125149 (2013); *Phys. Rev. Lett.*, **109**, 226401 (2012).
4. A. van Roekeghem et al., *Phys. Rev. Lett.*, **113**, 266403 (2014); *Europhysics Letters*, **108**, 57003 (2014).