

# Lecture 14: Multiple integrals and change of variables

Rafikul Alam  
Department of Mathematics  
IIT Guwahati

## Riemann sum for Triple integral

Consider the rectangular cube  $V := [a_1, b_1] \times [a_2, b_2] \times [a_3, b_3]$  and a bounded function  $f : V \rightarrow \mathbb{R}$ .

Let  $P$  be a partition of  $V$  into sub-cubes  $V_{ijk}$  and  $\mathbf{c}_{ijk} \in V_{ijk}$  for  $i = 1 : m, j = 1 : n, k = 1 : p$ . Also let

$\Delta V_{ijk} := \text{Volume}(V_{ijk}) = \Delta x_i \Delta y_j \Delta z_k$  and  $\mu(P) := \max_{ijk} \Delta V_{ijk}$ .

Consider the **Riemann sum**

$$S(P, f) := \sum_{i=1}^m \sum_{j=1}^n \sum_{k=1}^p f(\mathbf{c}_{ijk}) \Delta V_{ijk}.$$

## Triple integral

If  $\lim_{\mu(P) \rightarrow 0} S(P, f)$  exists then  $f$  is said to be **Riemann integrable** and the **(triple) integral** of  $f$  over  $V$  is given by

$$\iiint_V f(x, y, z) dV = \iiint_V f(x, y, z) dx dy dz = \lim_{\mu(P) \rightarrow 0} S(P, f).$$

**Theorem:** Let  $f : V \rightarrow \mathbb{R}$  is continuous. Then

- $f$  is Riemann integrable over  $V$ .
- **Fubini's theorem** holds, i.e, the iterated integrals exist and are equal to  $\iiint_V f dV$ .

## Example

Evaluate  $\iiint_V xyz^2 dV$  where  $V = [0, 1] \times [-1, 2] \times [0, 3]$ .

By Fubini's theorem,

$$\iiint_V f dV = \int_0^3 \left( \int_{-1}^2 \left( \int_0^1 x dx \right) y dy \right) z^2 dz = \frac{27}{4}.$$

## Triple integrals over general domains

Let  $D \subset \mathbb{R}^3$  be bounded and  $f : D \rightarrow \mathbb{R}$  be a bounded function. Then  $f$  is said to be **integrable over  $D$**  if for some rectangular cube  $V$  containing  $D$  the function

$$F(x, y, z) := \begin{cases} f(x, y, z) & \text{if } (x, y, z) \in D \\ 0 & \text{otherwise} \end{cases}$$

is Riemann integrable over  $V$ . Then

$$\iiint_D f(x, y, z) dV := \iiint_V F(x, y, z) dV$$

and

$$\text{Volume}(D) := \iiint_D dV.$$

## Type-I domain:

A domain  $V \subset \mathbb{R}^3$  is **Type-I** if

$$V = \{(x, y, z) : (x, y) \in D \text{ and } u_1(x, y) \leq z \leq u_2(x, y)\}$$

for some  $D \subset \mathbb{R}^2$  and **continuous functions**  $u_i : D \rightarrow \mathbb{R}$ .

If  $f : V \rightarrow \mathbb{R}$  be continuous and  $D$  is a **special domain** (e.g., Type-I, Type-II, Type-III) then

$$\iiint_V f(x, y, z) dV = \iint_D \left( \int_{u_1(x, y)}^{u_2(x, y)} f(x, y, z) dz \right) dx dy.$$

Similar results hold for **Type-II and Type-III domains**.




Figure: Type-I domain




Figure: Type-II domain






Figure: Type-III domain

## Example:

Evaluate  $\iiint_V 2x dV$  where  $V$  is the region bounded by the planes  $x = 0, y = 0, z = 0$  and  $2x + 3y + z = 6$ .

Note that  $V$  is Type-I:

$$0 \leq z \leq 6 - 2x - 3y \text{ and } (x, y) \in D,$$

where  $D$  is special domain given by

$$0 \leq x \leq 3 \text{ and } 0 \leq y \leq -\frac{3}{2}x + 2.$$

Thus

$$\begin{aligned} \iiint_V 2x dV &= \iint_D \left( \int_0^{6-2x-3y} dz \right) 2x dx dy \\ &= \int_0^3 \int_0^{-\frac{3}{2}x+2} (6 - 2x - 3y) 2x dx dy = 9. \end{aligned}$$

## Change of variable

Let  $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$  be  $C^1$  given by  $T(u, v) = (x(u, v), y(u, v))$ . Then the Jacobian matrix  $J(u, v)$  of  $T$  is given by

$$J(u, v) := \begin{bmatrix} x_u & x_v \\ y_u & y_v \end{bmatrix}.$$

Define the Jacobian of  $T$  by

$$\frac{\partial(x, y)}{\partial(u, v)} := x_u y_v - x_v y_u = \det J(u, v).$$

**Polar coordinates:** Define  $T(r, \theta) := (r \cos \theta, r \sin \theta)$ . Then

$$\frac{\partial(x, y)}{\partial(r, \theta)} = \begin{vmatrix} \cos \theta & -r \sin \theta \\ \sin \theta & r \cos \theta \end{vmatrix} = r.$$

## Change of variable for double integrals

Suppose  $T$  is injective and  $J(u, v)$  is nonsingular. Let  $D \subset \mathbb{R}^2$  and  $G := T(D)$ . Suppose that  $f$  is integrable on  $G$ . Then

$$dA = dx dy = \left| \frac{\partial(x, y)}{\partial(u, v)} \right| du dv$$

and

$$\iint_G f(x, y) dx dy = \iint_D f(x(u, v), y(u, v)) \left| \frac{\partial(x, y)}{\partial(u, v)} \right| du dv.$$

Polar coordinates:

$$\iint_G f(x, y) dx dy = \iint_D f(r \cos \theta, r \sin \theta) r dr d\theta.$$

## Example

Evaluate  $\iiint_G \sqrt{x^2 + z^2} dV$  where  $G$  is the region bounded by the paraboloid  $y = x^2 + z^2$  and  $y = 4$ .

We have

$$\iiint_G f(x, y, z) dV = \iint_D \left( \int_{x^2+z^2}^4 dy \right) f(x, y, z) dx dz,$$

where  $D = \{(x, z) : x^2 + z^2 \leq 4\}$ .

Setting  $x = r \cos \theta$  and  $z = r \sin \theta$  for  $(r, \theta) \in [0, 2] \times [0, 2\pi]$ ,

$$\iiint_G f(x, y, z) dV = \int_0^{2\pi} \int_0^2 r(4 - r^2) r dr d\theta = \frac{128\pi}{5}.$$

## Change of variable for multiple integrals

Let  $D \subset \mathbb{R}^n$  be open and bounded. Let  $T : D \rightarrow \mathbb{R}^n$  be such that  $T$  is  $C^1$ , injective and the Jacobian  $J(u)$  is nonsingular for  $u \in D$ .

Let  $G := T(D)$  and  $f : G \rightarrow \mathbb{R}$  be integrable over  $G$ . Then

$$dx_1 \cdots dx_n = \left| \frac{\partial(x_1, \dots, x_n)}{\partial(u_1, \dots, u_n)} \right| du_1 \cdots du_n$$

and

$$\begin{aligned} \int_G f(x) dx_1 \cdots dx_n &= \int_D f(x(u)) \left| \frac{\partial(x_1, \dots, x_n)}{\partial(u_1, \dots, u_n)} \right| du_1 \cdots du_n \\ &= \int_D f(x(u)) \left| \frac{dx}{du} \right| du. \end{aligned}$$

## Cylindrical coordinates

Consider  $T(r, \theta, z) = (r \cos \theta, r \sin \theta, z)$ . Then

$$\left| \frac{\partial(x, y, z)}{\partial(r, \theta, z)} \right| = \begin{vmatrix} \cos \theta & -r \sin \theta & 0 \\ \sin \theta & r \cos \theta & 0 \\ 0 & 0 & 1 \end{vmatrix} = r.$$

Thus  $dV = r dr d\theta dz$  and

$$\iiint_G f(x, y, z) dV = \iiint_D f(r \cos \theta, r \sin \theta, z) r dr d\theta dz.$$

## Example

Evaluate  $\iiint_G \sqrt{x^2 + y^2} dV$ , where  $G$  is the region bounded by  $x^2 + y^2 = 1$ ,  $z = 4$  and  $z = 1 - x^2 - y^2$ .

Consider cylindrical coordinates

$$D := \{(r, \theta, z) : (r, \theta) \in [0, 1] \times [0, 2\pi], 1 - r^2 \leq z \leq 4\}.$$

Then

$$\iiint_G f(x, y, z) dV = \int_0^1 \int_0^{2\pi} \left( \int_{1-r^2}^4 dz \right) r \, r dr d\theta = \frac{12\pi}{5}.$$



## Spherical coordinates

Consider  $T(r, \theta, \phi) = (r \sin \phi \cos \theta, r \sin \phi \sin \theta, r \cos \phi)$ .

Then

$$\begin{aligned} \left| \frac{\partial(x, y, z)}{\partial(r, \theta, \phi)} \right| &= \begin{vmatrix} \sin \phi \cos \theta & -r \sin \phi \sin \theta & r \cos \phi \cos \theta \\ \sin \phi \sin \theta & r \sin \phi \cos \theta & r \cos \phi \sin \theta \\ \cos \phi & 0 & -r \sin \phi \end{vmatrix} \\ &= -r^2 \sin \phi. \end{aligned}$$

Thus  $dV = r^2 \sin \phi dr d\theta d\phi$  and

$$\begin{aligned} \iiint_G f(x, y, z) dV &= \\ \iiint_D f(r \sin \phi \cos \theta, r \sin \phi \sin \theta, r \cos \phi) r^2 \sin \phi dr d\theta d\phi. \end{aligned}$$




Figure: Spherical coordinate system

## Example

Evaluate  $\iiint_G e^{x^2+y^2+z^2} dV$ , where  
 $G := \{(x, y, z) : x^2 + y^2 + z^2 \leq 1\}$ .

Using spherical coordinates we have

$$\begin{aligned} \iiint_D f(x, y, z) dV &= \int_0^\pi \int_0^{2\pi} \int_0^1 e^{r^2} r^2 \sin \phi dr d\theta d\phi \\ &= \frac{4}{3}\pi(e - 1). \end{aligned}$$

\*\*\* End \*\*\*